Forward-Central Jet Correlations at the Large Hadron Collider

Michal Deák

Instituto de Física Teórica,

Universidad Autonoma de Madrid

in collaboration with Francesco Hautmann, Hannes Jung, Krzysztof Kutak

(2010) 1012.6037

Contents

- Short introduction and motivation
- High energy factorisation framework
- High energy factorisation vs. multi parton interactions (MPI)
- Energy and jet flows
- Summary and conclusions

Forward jets - motivation

- QCD at small x

(Mueller, Navelet; Nucl.Phys. B282 (1987) 727)

- New particle discovery physics

- Extensive coverage of large rapidity regions at the LHC experiments ($3<|\eta|<5$ and $-5.2>\eta>-6.6$)
  - Possibility to study two jet correlations

Forward jets - relevant kinematics

- Equation for the rapidity of the hard subprocess final state

$$p = (p_0, p_3, p_2, p_1)$$

$$p_A = \left(\frac{\sqrt{s}}{2}, \frac{\sqrt{s}}{2}, 0, 0\right)$$

$$p_B = \left(-\frac{\sqrt{s}}{2}, \frac{\sqrt{s}}{2}, 0, 0\right)$$

$$y = \frac{1}{2}\ln\frac{p_0 + p_3}{p_0 - p_3} = \frac{1}{2}\ln\frac{x_1}{x_2}$$



Forward jets - relevant kinematics

- Equation for the rapidity of the hard subprocess final state

$$p = (p_0, p_3, p_2, p_1)$$

$$p_A = \left(\frac{\sqrt{s}}{2}, \frac{\sqrt{s}}{2}, 0, 0\right)$$

$$p_B = \left(-\frac{\sqrt{s}}{2}, \frac{\sqrt{s}}{2}, 0, 0\right)$$

$$y = \frac{1}{2}\ln\frac{p_0 + p_3}{p_0 - p_3} = \frac{1}{2}\ln\frac{x_1}{x_2}$$



- If $x_1 \sim 10^{-5}$ and $x_2 \sim 0.1$ then $y \sim 4.5$ -- very forward!

Simple kinematics + dynamics



- 2 distinct scales s_1 and s_2
- In the cross section of this process will appear logarithms of the form

$$\ln\left(\frac{s_2}{s_1}\right) \sim \ln\left(\frac{x_2}{x_1}\right) \overset{x_2 \simeq 1}{\sim} \ln\left(\frac{1}{x_1}\right)$$

- Can be resummed by high energy factorisation

  S. Catani, M. Ciafaloni and F. Hautmann, Phys. Lett. B242 (1990) 97; Nucl. Phys. B366 (1991) 135

- Exclusive states possible with **CCFM** equation implemented in Monte Carlo **CASCADE**
- Parton showers based on the **CCFM** equation

Cross section

- The dominant processes for forward jet production

$$qg^* \to qg$$
$$gg^* \to gg$$
$$gg^* \to q\bar{q}$$

- One parton off-shell carrying a small x
- Goal is to calculate the cross section

$$\frac{d\sigma}{dQ_t^2 d\varphi} = \sum_a \int \phi_{a/A} \otimes \frac{d\hat{\sigma}}{dQ_t^2 d\varphi} \otimes \phi_{g^*/B}$$

M. D., F. Hautmann, H. Jung, K. Kutak, JHEP **09**, 121 (2009). 0908.0538

- Off-shell matrix element convoluted with unintegrated parton density functions (uPDFs)
- In **CASCADE** done by generating emissions in a parton shower

Matrix element study

- Some properties of the off-shell matrix elements of the relevant processes studied - decomposed into non-abelian and abelian part





$$qg^* \longrightarrow qg$$
$$gg^* \longrightarrow gg$$
$$gg^* \longrightarrow q\bar{q}$$

Q_T=transversal energy in terms of final state pt's

- Dependence on transvrsal momentum of the off-shell gluon
- Dynamical cut-off at $k_T \sim Q_T$ set by coherence effects
  - Non-negligible contribution from finite tail
- DGLAP based parton showers do not allow for such a hard emissions in the chain

M. D., F. Hautmann, H. Jung, K. Kutak, JHEP **09**, 121 (2009). 0908.0538

CCFM equation

- The CCFM equation
  - Includes BFKL kernel
  - Coherence effects
  - Angular ordering
- The equation



$$\mathcal{F}(x,\mathbf{k},\mathbf{q}'^2) = \mathcal{F}(x,\mathbf{k},\mathbf{q}_0'^2) + \int_{\mathbf{q}_0'^2}^{\mathbf{q}'^2} \frac{d^2\bar{\mathbf{q}}'}{\bar{\mathbf{q}}'^2} \frac{N_C\alpha_S}{\pi}$$

$$\int_x^{1-\frac{Q_0}{|\mathbf{q}'|}} \frac{dz}{z} \mathcal{F}(x/z,\mathbf{k}',\bar{\mathbf{q}}'^2) \left(\frac{\Delta_{NS}(\mathbf{k}'^2,(z\bar{\mathbf{q}}')^2)}{z} + \frac{1}{1-z} \right) \Delta_S(\mathbf{q}_0'^2,(z\bar{\mathbf{q}}')^2)$$

M. Ciafaloni, Nucl. Phys. B296, 49 (1988);
S. Catani, F. Fiorani, and G. Marchesini, Phys. Lett. B234, 339 (1990);
S. Catani, F. Fiorani, and G. Marchesini, Nucl. Phys. B336, 18 (1990);
G. Marchesini, Nucl. Phys. B445, 49 (1995)

CASCADE

- Monte Carlo generator **CASCADE** (version 2.2.0, H. Jung et al.; Eur.Phys.J.C70:1237-1249,2010) - implementation of the CCFM equation
  - Backward evolution algorithm for initial state parton showers for
    - Exact kinematics in each step of the parton shower
    - No difference between parton shower evolved uPDF and CCFM evolved uPDF
    - Gluon chains
    - Valence quarks/Non-singlet uPDFs from one-loop CCFM equation
  - Final state parton showers by Pythia algorithm
  - Hadronisation of partons by the Lund String Model
  - Gluon uPDFs obtained from fits to HERA data

Details of the convolution

- The initial state off-shell gluon directly from the CCFM-parton shower-evolved uPDF

$$qg^* \longrightarrow qg$$
$$gg^* \longrightarrow gg$$
$$gg^* \longrightarrow q\bar{q}$$

- On-shell parton obtained from CCFM evolved uPDF
  - Transversal momentum neglected in the matrix element of the hard subprocess, but included in the kinematics of the final state
  - $k_\perp$ integrated up to the hard scale given by the angle

- Quarks evolved by one-loop CCFM: only valence component included

H. Jung et al.; Eur.Phys.J.C70:1237-1249,2010



High energy factorisation and MPI

- 1. MPI effectively generate certain amount of pt-non-ordering of showered partons
- 2. MPI increase the jet multiplicity of the parton showers
- 3. MPI increase the number of hard jets in the process
- pt-unordered parton showers (1.) increase jet Multiplicity (2.) and number of hard jets (3.)

- **Natural to compare MPI with pt-unordered parton Showers**
- Different mechanisms



multiple pt-ordered chains

one pt-unordered chain

Phenomenological results

Parameters and settings

- Hard matrix elements included in Cascade MC generator:
  - gluon uPDFs fitted from HERA data
  - quark CTEQ6.0 PDF evolved by CCFM
- Pythia settings: - CTEQ 5L PDFs
  - q^2 ordering for initial state parton shower, with and without MPI
- Jets on parton level; jet algorithm - kt-clus

- Forward jets rapidity: $-3 < \eta < -5$
- Central jets rapidity: $-2 < \eta < 2$
- $E_\perp > 10$ GeV

Pt spectra of forward jets

Forward jet
$-3 < \eta < -5$





- Similar slopes of cross sections
- MPI only shift the jet rapidity cross section by a factor

Pt spectra of central jets

Central jet
$-2 < \eta < 2$





- Different slopes of cross sections
- k_T of incoming gluon allows for harder spectrum - CCFM parton showers not ordered in k_T
- MPI enhances cross section at low E_T

Rapidity dependence





- MPI changes the slope of the rapidity spectrum
- Rapidity cross sections agree better when increasing the cut on jet E_T

Azimuthal dependence



- flattening dependent on $\Delta\eta$

Forward jets - Summary

- Calculations of gauge invariant matrix elements of relevant processes in Monte Carlo implementation suitable form
- Convolution with the uPDFs in Monte Carlo generator **CASCADE**
- Study of jet transversal energy and rapidity cross sections of forward jets and a central jet
- Comparison with **PYTHIA** Monte Carlo generator with multi-parton interactions (MPI)
- Difference between the high energy factorisation enhanced jet activity and MPI jet activity

Conclusions

- LHC opens phase space for large center of mass energies and for presence of multiscales

- This brings perturbative corrections which are summed up by high energy factorisation

- An approach which allows for studies of forward jets at the LHC

- Proposal of observables which allow for discrimination between different approaches

CASCADE

- The largest angle = the angle of the hard subprocess final state system

1.



2.



...

- Angular ordering for small angles

$$\frac{|\mathbf{q}_i|}{1-z_i} > \frac{z_{i-1}|\mathbf{q}_{i-1}|}{1-z_{i-1}}$$