QCD results from the Tevatron

Dmitry Bandurin

Florida State University

On behalf of D0 and CDF Collaborations

DIS 2011, April 11, Newport News, VA, USA

Tevatron collider



Tevatron collider luminosity



- D0 and CDF recorded about $10\,fb^{-1}$
- Efficiency of data taking >90%
- Current results correspond to luminosity of $1-6\,fb^{-1}$

Outline

- Jet production:
  - Inclusive jets
  - Dijets
  - 3-jets
- V(=W,Z) + jets production
  - V + inclusive jets
  - V + heavy flavor jets
- Inclusive photon and di-photon production
- Underlying events and Double parton interactions



Jet Results

PDF, αs

Searches for New Physics

Motivations for the jet measurements

- Well-understood by NLO pQCD
- PDF constrain

– x-Q^2 regions accessible at fixed target, DIS, Tevatron and LHC are complementary to each other

– only Tevatron incl. jet data provide significant constraint on gluon PDF at high x and high Q^2

- New Phenomena searches:

– particles decaying to jets, ED, quark compositeness, etc

– searches for new phenomena are limited without proper understanding QCD background





Inclusive jet production (CDF)

Inclusive jet measurements test pQCD over 8 orders of magnitude in 5 rapidity regions up to jet pT ~600 GeV.

- CDF measured inclusive jet cross section with Midpoint cone algorithm (R=0.7) and kT (D=0.4, 0.7, 1.0) algorithm.
- Data/Theory consistent for the cone and kT (for all D parameters) algorithms
  => both algorithms can be successfully used at hadron colliders.

PRD78, 052006 (2008)



PRD75, 092006 (2007)



Inclusive jet production (D0)

PRL 101, 062001 (2008)

D0 also measured inclusive jet cross section using Midpoint algorithm in 6 rapidity regions.

Dominant systematic uncertainty is from JES:
Steeply falling spectrum:
=> Even small JES uncertainty leads to large uncertainties on cross section

Typical JES uncertainty:
1-2% in D0, 2-3% in CDF

Total uncertainty on the cross sections:
15-30% in D0, 15-50% in CDF





Energy scale uncertainty, D0

Inclusive jet production: Data/Theory (D0,CDF)

In general, CDF and D0 measurements are in agreement with QCD NLO predictions.

However, data favored lower bound of the theoretical (CTEQ6.5M PDF) predictions, with smaller gluon content at high x.

Experimental uncertainties at high pT are lower than theoretical (largely PDF ones):
=> constrain PDF





Leads to modified central values (esp. at x>0.3) and reduced PDF uncertainties.
(see also p.50 in Backup)

D0 results are most precise measurement to date.

MSTW 2008 uses CDF kT and D0 cone results.

Inclusive jet production: hadron colliders





pT plot: the Tevatron pT reach is still about as good as the published LHC results
xT plot: the Tevatron data have far better high-x sensitivity

Measurement of α_s from inclusive jets (D0)

PRD 80, 111107 (2009)

- Cross section formula:

$$\sigma_{\text{theory}}(\alpha_s) = \left(\sum_n \alpha_s^n c_n \right) \otimes f_1 \otimes f_2$$



- c_n: perturbative coefficients (→ pQCD matrix elements)
- f_1, f_2: PDFs of colliding $p, \bar{p}$

Determine α_s from data:

- Vary α_s until σ_{theory} agrees with σ_{exper}
- ...for each single bin →
- χ^2 fit of theory to data (p. 62 in backup) using 21 NNLO PDF sets from MSTW2008 with α_s within 0.107-0.127 in 0.001 steps
- 5 NLO CTEQ6.6M sets are also considered
- Only 22 points of 110 are used (with x<0.2)



Running of $\alpha_s(pT)$

- Combine points in different |y| regions at same pT
  → Produce 9 $\alpha_s(pT)$ points from selected 22 data points



theory:NLO+2-loop threshold corrections

→ About same precision as HERA jets (**0.1189 ±0.0032**)

→ The only Run II result on α_s

→ Improvement as comp. with Run I
(**0.1178±0.0001**(stat)$^{+0.0081}_{-0.0095}$(syst))

Compare to HERA results:

→ consistency

→ extend pT reach of HERA results to higher pT range of 50-145 GeV

"World average": **0.1184±0.0007**

PRD 80, 111107 (2009)

Dijet mass cross section measurement (D0)

PLB 693, 531 (2010)



- Measurement of dijet mass in six rapidity bins, $|y|_{max} = \max(|y_1|, |y_2|)$

Non-perturbative corrections (-10%, 23%)

Comparison to NLO pQCD with MSTW2008 and CTEQ6.6M NLO PDFs, $\mu_F = \mu_R = (pT_1 + pT_2)/2$



– 40—60% difference between PDFs (MSTW2008/CTEQ6.6) at high masses

– Data/QCD in good agreement in central region

– Data are lower than central pQCD prediction at higher rapidities

Last mass bin is at ~1.3 TeV!

Dijet mass cross section measurement (CDF)

PRD 79, 112002 (2009)



Study dijet events in |y|<1.0

(uses same dataset as the inclusive jets)

=>New physics expected to be produced more centrally & expect better S/B in central region

Total uncertainty: $^{+13}_{-12}$% at low m_{jj}

$^{+76}_{-49}$% at high m_{jj}

– NLO pQCD fits to data: χ^2/ndf = 21/21

(syst. uncertainties and non-perturbative corrections all independent; fully correlated over m_{jj})

– Data/QCD agreement similar to D0 for the central region

PARTON-TO-HADRON LEVEL CORRECTION

Pythia (TuneA) central value; Herwig PS taken as uncertainty



Dijet mass: searches for new physics (CDF)

PRD 79, 112002 (2009)

Dijet mass tests pQCD but also sensitive to presence of new physics, resonances decaying to two jets

=> Use uncorrected jet data to maximize sensitivity to resonances

No significant evidence for resonant structure has been observed, so set limits



Observed mass exclusion range	Model description
260-870 GeV/c^2	Excited quark → qg (f=f'=f$_s$=1)
260-1100 GeV/c^2	ρ_{T8} techni-rho
260-1250 GeV/c^2	Axigluon/coloron
290-630 GeV/c^2	E_6 diquark
280-840 GeV/c^2	W' (SM couplings)
320-740 GeV/c^2	Z' (SM couplings)

D0 dijet χ: limits on q-compositeness, Extra Dim.: PRL 103, 191803 (2009)

Three jet mass cross section (D0)

Submitted to Phys.Lett.B

Differential measurements of 3-jet mass:

- p_T^{lead}>150 GeV, p_T^{3rd} >40 GeV; ΔR_{jj}>1.4

– Measurement is done in 3 rapidity and 3 pT intervals of 3rd jet.

– Three-jet calculation available @NLO

- Used fastNLO with MSTW2008
- Default scale mu = 1/3(pT1+pT2+pT3)
- Scale uncertainties: independent variations by *x*2 of renorm. and factor. scales

– NLO non-perturbative corr.: -3%,+6% (DW used as a default, x-checked with tunes A,BW, Z1,Perugia soft&hard)

– Total systematic uncertainty: 20-30% (dominated by JES, p_T resolution and lumi)



Three jet mass cross section (D0)

Differential in y

Differential in p_T



χ^2 test for *central* PDF values

TABLE II: χ^2 values between data and theory for different PDF parametrizations in the order of decreasing χ^2, for all 49 data points.

PDF set	Default $\alpha_s(M_Z)$	χ^2 at $\mu_r = \mu_f = \mu_0$ for default $\alpha_s(M_Z)$	χ^2_{minimum}
HERAPDFv1.0	0.1176	95.1	81.7
CT10	0.1180	94.5	88.2
ABKM09NLO	0.1179	76.5	76.5
NNPDFv2.1	0.1190	65.9	63.3
MSTW2008NLO	0.1202	59.5	59.5

- Good agreement seen between data and NLO (MSTW2008) for all cases.
- Comparisons to ABKM09, NNPDF2.1, HERA1.0 are also provided.
- χ^2 test is done for 3 theor. scales and all α_s values available for a given PDF set
- Best χ^2 results for MSTW2008, NNPDF2.1

See also talks by Z.Hubacek & M.Wobisch on April 12 at QCD & Had F.S.

Ratio of 3 to 2 jet production cross sections (D0)

Preliminary

- First measurement of ratios of multijet cross-sections at Tevatron
- Test of QCD almost independent of PDFs
- Many experimental uncertainties also cancel in the ratio R3/2.
- Measure $R_{3/2}$= P(3rd jet | 2 jets) as a function of two momenta p_{Tmax}, p_{Tmin}:
  - p_{Tmax} – leading jet p_T (common between 2- and 3-jet productions)
  - p_{Tmin} – scale at which other 1-2 jets resolved
- Comparisons to NLO QCD, LO Sherpa and Pythia with a few tunes

- Shape of the ratios is well described by NLO theory and, as expected, practically independent on PDF set.
- Excellent agreement to Sherpa 1.1.3 (MSTW2008 LO), Pythia BW tune (tunes QW, DW [they worked for χ^2, $\Delta\varphi$ data], Perugia are significantly off)
- Probes running of α_s up to p_T of 500 GeV



Structure of high pT jets (CDF)

Preliminary

- Motivation: (a) test of QCD, tuning parton showering mechanism
  (b) can be used for new physics searches with
  a heavy resonance decay (Higgs, neutralinos, high pT top-quarks)

- Mass is calculated using standard E-scheme: 4-vector sum over towers in a jet, which gives (E,px,py,pz)
- Selections: ≥1 jet with pT>400 GeV, 0.1<|y|<0.7: 3136 (3621) events, jet R=0.4-1.0
  anti-top: m_jet2<100 GeV and S_met < 4 and pT_jet2>100 GeV



400<pT<500 GeV, anti-top cuts



Mass of high pT jets: comparison with theory (CDF)

– Good agreement between data and QCD LLA and Pythia predictions over jet mass range 100-250 GeV and for both jet cones, R=0.4 and 0.7.

– Data interpolate between QCD predictions for quark and gluon jets; about 80% of high mass jets are caused by quark fragmentation.
(See 0807.0234, 0810.0934 for more on the theory)

– Jet angularity and planar flows have also been studied (see backup slides 59,60): Data prefer more "spherical" and aplanar configurations than QCD predictions.

R = 0.4



R = 0.7





V + jet Results

Fixed-order: NLO

LO + Parton Shower

Backgrounds to New Physics

Vector Boson + Jet

WH production



Single top production



W+dijet, M_{jj} "bump"



– Background to top-quark, Higgs, SUSY, other NP productions

– Provide detailed measurements of p_T and angular distributions of vector boson and jet

→ test of fixed order perturbative QCD, LO ME+PS predictions in EvGen

→ testing and tuning of phenomenological models

Z+jets production: jet pT, #jets (CDF)

Preliminary

6 fb-1, ee and $\mu\mu$ channels, jet pT>30 GeV and |y|<2.1





- Good agreement with QCD NLO in jet pT and #jets.

Z+jets production: jet pT, data/NLO (D0)

Measurement of 1st, 2nd and 3rd jet pT in Z events:

PLB 678, 45 (2009)

- Z→ee, jet p_T > 20 GeV, jet |y|< 2.5.
- Normalized to inclusive Z production x-section (cancel some uncertainties)

1st jet in Z + jet + X | **2nd jet in Z + 2jet + X** | **3rd jet in Z + 3jet + X**



Z+jets production; jet pT, data/MC (D0)

Comparison to Pythia, Herwig, Alpgen and Sherpa
Treating the scale choice as a tuneable parameter:
best description from Alpgen with lower scale (default: $\mu_F^2 = p_{TZ}^2 + M_Z^2$).

Leading jet in Z + jet + X **2nd jet in Z + 2jet + X** **3rd jet in Z + 3jet + X**



W+jets production (CDF)

Preliminary

- Jets are defined with MidPoint R=0.4, $p_T \geq 20$ GeV/c, $|\eta| \leq 2.0$
- $W \rightarrow e\nu(\mu\nu)$, $|\eta_{e(\mu)}| \leq 1.1$, $p_T[e(\mu)] \geq 20$ GeV, $M_T[W] \geq 40(30)$ GeV for $e(\mu)$





- Data are compared to Alpgen+Pythia
- For the pT spectra MC are normalized to total x-section in data with ≥1 jet
- Good agreement data/MC for jet pT and #jets.

See also talks on W+jets production at CDF and D0 at QCD & Had.F.S. (April 14)

σ(Z+b) / σ(Z+jet) (D0)

- Important background to the SM Higgs search in the ZH channel.
- Probe of b-quark PDF, important for gb → Hb & single-top studies
- Measurement of ratio σ(Z+b) / σ(Z+j) benefits from cancellation of many systematics => precise comparison with theory



- L = 4.2 fb-1
- Z → ee/μμ + b + X
  $70 < M_Z < 110$ GeV
- lepton $p_T > 15$ GeV
- D0 RunII Midpoint Cone jets with R=0.5
- jet $p_T > 20$ GeV
- jet $|\eta| < 2.5$

- Secondary vertex tagging
  → Apply Neural Network algorithm on jets to enrich data with b-jets (NNout > 0.5)
  → Use a longer b-hadron lifetime to discriminate between b/c/light jets
- Use data for light jet template, Pythya+Alpgen for b & c jets
- Use log likelihood fit to extract b- jets fractions



σ(Z+b) / σ(Z+jet) (D0)

PRD83, 031105 (2011)





- Measurement: **0.0193 ± 0.0022 (stat)± 0.0015 (syst)** [~8% syst]
  Most precise measurement of 'Z+b' fraction to date!
- Consistent with NLO theory**: 0.0192 +/- 0.0022**
  (MCFM, renorm. and factor. scales are at Mz)

Z+b production (CDF)

PRD79, 052008 (2009)

- L=2 fb-1
- Z → ee/μμ + b + X
- jet p_T > 20 GeV, jet |η| < 1.5
- Jet track mass in the secondary vertex is used to discriminate between jet flavors

Theory:

- MCFM : all calculations are at O(αs^2)
- Pythia, Alpgen



σ(Z+b) / σ(Z)

Data: [3.32± 0.53(stat)±0.42(syst)]$x10^{-3}$

MCFM: 2.3 (2.8) $x10^{-3}, Q^2 = Mz^2(jet\ p_T^2)$

Pythia: 3.5 $x10^{-3}$

Alpgen: 2.1 $x10^{-3}$



σ(Z+b) / σ(Z+jet)

Data: 2.08 ± 0.33(stat)±0.34(syst) %

MCFM: 1.8% / 2.2%

Pythia: 2.2%

Alpgen: 1.5%

MCFM(μ= jet p_T^2), Pythia provide best agreement to data

σ(W+b) (CDF)

L= 1.9 fb-1

Phys.Rev.Lett.10,131801 (2010)

$p_T^{b\text{-jet}} > 20$ GeV, $|\eta_{b\text{-jet}}| < 2.0$, ($R_{\text{cone}} = 0.4$), $p_T^{\ell} > 20$ GeV, $|\eta_\ell| < 1.1$, $p_T^V > 25$ GeV



- $\sigma(W_{(\to \ell\nu)} + b) = \dfrac{N_{b\text{-tags}} \cdot f_{b\text{-jets}} - N_{\text{bkg}}^{b\text{-jets}}}{\mathcal{L} \times A \times \epsilon}$

Major *b*-jet backgrounds:

$t\bar{t}$ (40% of total background)
single top (30%)
Fake W (15%)
WZ (5%)

Measured cross section larger than ALPGEN and NLO predictions ($\sim 3\sigma$)

σ(W+b-jets) · BR(W → l ν) = 2.74 ± 0.27 (stat) ± 0.42(syst) pb

QCD NLO: 1.2 + 0.14 pb
Alpgen : 0.78 pb

About a factor 2(3) of descrepancy with NLO (Alpgen)!

Photon production

Test fixed order NLO, resummation, fragmentation, PDF

Photon Production



Direct photons emerge unaltered from the hard subprocess
→ direct probe of the hard scattering dynamics
→ potential sensitivity to PDFs (gluon!)
...but only if theory works



also fragmentation contributions:



suppress by isolation criterion
→ observable: **isolated** photons

32

Inclusive Isolated Photons (CDF,D0)

Phys.Rev.D80,111106 (2009)



Phys. Lett. B 639, 151 (2006)



CDF and D0: 20< p_T <400 GeV, both results are for central rapidities, *same binning*

- D0/CDF: results in agreement
- Data/Theory: difference in low p_T shape
- experimental and theory uncertainties > PDF uncertainty
  → no PDF sensitivity yet
- First: need to understand discrepancies in shape (similar to results of UA2, CDF Run 1)

Photon Pair Production (D0)

- Almost irreducible background to H→ $\gamma\gamma$, other new phenomena, => should be understood
- Isolated (ETsum[R=0.4]< 2.5 GeV) photons with pT>20 and 21 GeV, |y|<0.9; 4.2 fb-1
- Data are compared with predictions by PYTHIA, DiPhoX, ResBos
- 1D cross sections in diphoton Mass, $p_T^{\gamma\gamma}$, $\Delta\phi$, cosθ* and 2D ones ($p_T^{\gamma\gamma}$, $\Delta\phi$, cosθ* in Mass bins)

Phys.Lett. B690,108(2010)





Good agreement between data and RESBOS for M>50-60 GeV

Data $p_T^{\gamma\gamma}$ spectrum is harder than predicted: need for NNLO? Unaccounted fragm. contribution?

Photon Pair Production (CDF)

http://www-cdf.fnal.gov/physics/new/qcd/diphoXsec_2010/public_diphoton.html

– Isolated (ETsum[R=0.4]< 2 GeV) photons with pT>15 and 17 GeV, |y|<1.0; 5.4 fb-1

– Data are compared with predictions by PYTHIA, DiPhoX, ResBos

– 1D cross sections vs. diphoton Mass, $p_T^{\gamma\gamma}$, $\Delta\phi$.

Preliminary





– None of the models describe the data well in all kinematic regions, in particular at low diphoton mass (M<60 GeV), low $\Delta\phi$ (<1.7 rad) and moderate $p_T^{\gamma\gamma}$ (20-50 GeV)

– Data/Theory: similar conclusion to those from D0 results

– See also a talk on recent D0 diphoton results on April 14 at QCD & Had.F.S.



Underlying events
Double parton scattering

Tuning phenomenological models, MC Generators

Underlying Event in DY and Jet production (CDF)

UE events: MPI + beam remnants
Goal: improve understanding and modeling of high energy collider events

Define 3 regions in a jet/DY event, based on the leading jet/dilepton pT
"toward"
"away"
"transverse"





"transverse" region
→ very sensitive to underlying event

Study (in all regions)

- charged particle density (per $\Delta\eta\Delta\phi$)
- multiplicity
- p_T sum density

Underlying Event in DY and Jet production(CDF)

Phys.Rev.D82,034001 (2010)

Comparison of **three regions** in DY:

- "away" region: pT density increases with lepton pair (or jet) pT
- "transverse", "toward" regions: pT density is almost flat with lepton pair pT





Comparison of "transverse" region between **jets and DY**

- similar trend in both (MPI universality?)
- tuned PYTHIA (A,AW) describes data

Minimum bias track multiplicities and pT (CDF)

→ Sensitive distribution to QCD perturbative/non-perturbative effects, MPI.

Phys.Rev.D79,112005 (2009)





→ Well described by "Tune A" MPI model

→ Not all processes are included in Pythia "minbias" event: source of a discrepancy at high track pT.

→ Data favors the presence of multiple parton interactions (MPI) and can be used to constrain MPI models. (MPI lead to larger Nch that are harder than the beam remnants but not as hard in pT as for the primary hard 2→2 scattering.)

Double Parton Scattering in γ+3 jet events (D0)

- Study of MPI events in high pT regime (jet pT>15 GeV); complementary to CDF.
- Complementary information about proton structure: Spatial distribution of partons
  ⇒ Possible parton-parton correlations. Impact on PDFs?
- Needed for understanding multijet signal events and correct estimating backgrounds to many rare processes.

Selections: 60 < photon pT < 80 GeV, lead. jet pT > 25, other 2 jets with pT > 15 GeV



Main Background: Single Parton scattering



$$\sigma_{DP} = \sigma_{\gamma j}\, \sigma_{jj}/\sigma_{eff}$$

Three types of events with Double Parton scattering



σ_{eff} is a scale parameter sensitive to the size of effective parton interaction region, and thus
=> to the parton spatial density

Double parton results, γ+3 jet events (D0)

Phys.Rev.D81,052012 (2010)





- The measured double parton fraction drops from 0.47±0.04 at $15<p_{T2}<20$ GeV to 0.23±0.03 at $25<p_{T2}<30$ GeV
- Effective cross section averaged over 3 p_{T2} bins:

$$\sigma_{eff}^{ave} = 16.4 \pm 0.3(stat) \pm 2.3(syst)\, mb$$

- Good agreement with Run I measurements by CDF ("4 jets", $\sigma_{eff} = 12.1^{+10.7}_{-5.4}$ mb and "γ+3jets", $\sigma_{eff} = 14.5 \pm 1.7^{+1.7}_{-2.3}$ mb)

Angular decorrelations in γ+2(3) jet events (D0)

Motivations:

- By measuring **differential** cross sections vs. azimuthal angles in γ+3(2) jet events we can better tune (or even exclude some) MPI models in events with high pT jets.

- Differentiation in jet pT increases sensitivity to the models even further.

Four normalized differential cross sections are measured

- $\Delta\phi(\gamma$+jet1, jet2) in 3 bins of 2nd jet pT: 15-20, 20-25 and 25-30 GeV
- $\Delta S(\gamma$+jet1, jet2+jet3) for 2nd jet pT 15-30 GeV (larger for stat. reasons but still has good sensitivity to MPI models)





See also talk at QCD & Had.F.S. on the double parton production at D0 (April 14)

$\triangle S$ and $\Delta\phi$ cross sections

Phys.Rev.D83,052008 (2011)





- MPI models substantially differ from any SP (=single parton scattering) prediction.
- Large difference between SP models and data confirm presence of DP events in the data sample.
- MPI models differ noticeably between each other, especially at small azimuthal angles
  => we can tune the MPI models or just choose the best one(s)
- Data are close to `Perugia (P0), S0 and Sherpa with MPI tunes.`
  N.B.: the conclusion is valid for both the considered variables and 3 jet pT intervals!

Summary

- A few recent Tevatron results are presented: current level of understanding jet ID, systematics and jet energy scale leads in many cases to experimental uncertainties similar or lower than theory uncertainties.
=> Precision measurement of fundamental observables.

- Good consistency between D0 and CDF in most cases, complementarity.

- Jet results: good agreement with pQCD, sensitivity to PDF sets, strongest constraint on gluon PDF, extraction of α_s, detailed studies of the effect of different jet algorithms; jet substructure, limits on many NP models.

- Z/W results: extensive tests of pQCD and MC models

- Photon results: test fixed order NLO, resummation, fragmentation. Theory should be better understood.

- Underlying/DP events: strong constraints/improving phenomenological models at low and high pT regimes.

Tevatron talks in parallel sessions

QCD and Hadronic Final States

Title	Speaker	Date
• Multi-jet measurements at D0	Z. Hubacek	Tuesday, April 12
• Jet production and the determination of the strong coupling constant at D0	M. Wobisch	Tuesday, April 12
• Light resonances in minimum bias events at CDF	S.Oh	Wednesday, April 13
• Direct Photon Pair Production at D0	D.Bandurin	Thursday, April 14
• Multi-parton interactions in photon+jets events at D0	A.Verkheev	Thursday, April 14
• W/Z + jets at CDF	D.Stentz	Thursday, April 14
• W/Z + jets at D0	D.Price	Thursday, April 14

Small-x, Diffraction and Vector Mesons

• Diffraction results from CDF	D.Goulianos	Wednesday, April 13

BACK-UP SLIDES

The DØ and CDF detectors

- Data taking efficiency (DØ & CDF) $\gtrsim$ 90%



- Multi purpose detectors with broad particle ID capabilities
- Stable detectors and triggers



- Calorimeters ($\rightarrow$ jets, e, γ): Fine granularity and good energy resolution
  DØ: $\Delta\eta \times \Delta\phi \sim 0.1 \times 0.1$
  CDF: $\Delta\eta \times \Delta\phi \sim 0.1 \times 0.26$
- Central tracking systems ($\rightarrow$ charged particles)
- Muon spectrometers ($\rightarrow$ muons)

Corrections to particle level



Data

Theory

In Run II jet results, in most cases:

- data are corrected to particle level
- particle level measurements are compared to NLO theory
- NLO theory is corrected to particle level using parton shower MC

$$C_{had} = \frac{\text{observable (particle level)}}{\text{observable (parton level)}}$$

- There is also correction (C_{ue}) for the underlying events (MPI). Usually we run Pythia with a couple of Tunes, Herwig+Jimmy and correct predictions with MPI to that without.

Inclusive Jets: Tevatron vs. LHC



PDF sensitivity:

→ compare jet cross section at fixed $x_T = 2\,p_T$ / sqrt(s)

Tevatron (ppbar)

>100x higher cross section @ all x_T

>200x higher cross section @ x_T>0.5

LHC (pp)

- need more than 2400 fb^{-1} luminosity to improve Tevatron@12 fb^{-1}
- more high-x gluon contributions
- but more steeply falling cross sect. at highest p_T (=larger uncertainties)

→ Tevatron results will dominate high-x gluon for some years

Gluon PDF and Tevatron data

from MSTW2008 paper



–CTEQ6.6 does not use Tevatron Run II jet data, while MSTW does

– MSTW2008 and CTEQ6.6 results are in agreement for x<0.3

=> Tevatron jets mostly affect PDF at x>0.3

Difference between quark and gluon responses

Responses in the calorimeter for quark and gluon jets are different
=> Different corrections are need depending on final state
(dijet events are dominated by the gluon jets, ttbar ones are quark dominated, etc)



JCCA – midpoint cone R=0.7

Jet energy scale calibration

- We do not "see" partons or particles in calorimeter, only ADC counts
- ADC counts --> cell energies
- Run jet cone algorithm (see Backup) with
  $\Delta R = \sqrt{(\Delta y^2 + \Delta\Phi^2)} < R_{cone}$

Jet's E are corrected to the particle level using the Jet Energy Scale (JES) setting procedure :

- Calibrate using γ+jets (dijets and Z+jets)
- JES includes: Energy Offset (energy not from the main hard scattering process); Detector Response, Out-of-Cone showering; Resolution
- Responses in the calorimeter for quark and gluon jets are different: additional corrections are applied to convert γ+jet → dijet JES.

Energy scale uncertainty: 1-2.5% (a lot of hard work of many people)!





Inclusive jet production (D0): correlations study

- All systematic uncertainties in data compose 24 main groups
- Possibility to constrain PDF further using the provided correlation matrices
- Detailed paper on the measurement to be submitted soon to PRD



Main sources of systematic uncertainties

Photon Pair Production (CDF)

Direct Photon Pair Production Differential Cross Sections measured with the CDF Detector: Ratios of Data/Theories





Resummation

Resummation

Fragmentations

Z/W+jets production

Use leptonic Z/W decays as most precise probe of QCD
- high Q^2 (~M_Z or M_W)
- very small backgrounds, right down to very small p_T!



Concentrate on high pT final states
- regime of perturbative QCD

Theory predictions:

pQCD (+ corrections for underlying event & hadronization):
- LO Z(W) + 1 - 6 partons
- NLO Z(W) + 1, 2 (MCFM)

[NLO W+3 (Rocket, Blackhat+SHERPA) is also available now]



Event generators:
- LO 2 ->1, 2 + parton shower
  - PYTHIA, HERWIG
- LO 2 -> 1-6 + (vetoed) parton shower
  - ALPGEN (MLM ME-PS matching),
  - SHERPA (CKKW ME-PS matching)



These generators are the main Tevatron and LHC tools
- but, leading order → large uncertainties
- must to be tuned to data!

σ(W+c)/σ(W+jet) at D0 and σ(W+c) at CDF

Sensitive to s-quark PDF: 90% s, 10% d

signal: OS>>SS
backgrounds: OS~SS
(subtracted in the diff. 'OS-SS')

Measurement cuts:
lepton p_T > 20 GeV
missing E_T > 20 GeV
D0 midpoint jet R_{cone}=0.5,
p_T^{jet} > 20 GeV,$|\eta^{jet}|$ < 2.5(1.5)



Phys.Lett.B666, 23 (2008)



D0 Data: 0.074 ± 0.019 (stat) ± $^{+0.012}_{-0.014}$ (sys)
Alpgen+Pythia: 0.044±0.003

CDF: σ(W+c)*Br(W-> lnu), L=1.8 fb-1 :
CDF Data: 9.8±3.2 pb
QCD NLO: 11.0 +1.4/ -3.0 pb

Good agreement data/theory

Phys.Rev.Lett.100,091803 (2008)

Angularity and planar flow (CDF)

- Angularity and planar flow variables study the jet substructure; quite robust against soft radiation, less dependent on the jet algorithm used.

– Angularity: sum over calorimeter towers:

$$\tau_a(R, p_T) = \frac{1}{m_J} \sum_{i \in jet} \omega_i \sin^a \theta_i \left[1 - \cos\theta_i\right]^{1-a} \sim \frac{2^{a-1}}{m_J} \sum_{i \in jet} \omega_i \theta_i^{2-a}$$

where ω_i is energy of a jet tower (particle)

– It is sensitive to the degree of symmetry in the energy deposition inside a jet: can distinguish jet originating from regular QCD production of light quarks and e.g. gluons from boosted heavy particle decay.

– Data show fewer jets at lower angularity, i.e. prefer more 'spherical' jets.



Angularity and planar flow (CDF)

– Planar flow is another jet substructure variable:

$$I_w^{kl} = \frac{1}{m_J}\sum_i w_i \frac{p_{i,k}}{w_i}\frac{p_{i,l}}{w_i} \qquad Pf = 4\frac{\det(\mathrm{I_w})}{\mathrm{tr}(\mathrm{I_w})^2} = \frac{4\lambda_1\lambda_2}{(\lambda_1+\lambda_2)^2}$$

where w_i is energy of a jet tower (particle), p_i,k is a k-th component of transverse momentum relative to the jet momentum axis; λ_1,2 is eigenvalue of the matrix I_w.

– ***Pf*** should vanish for linear shapes and close to unity for isotropic depositions of energy.

– At high jet masses (140-200 is considered) data prefer more aplanar configuration than QCD prediction (anti-top cuts are applied).





Combined $\alpha_s(M_Z)$

Based on 22 inclusive jet data points with x-test<0.15

Combined $\alpha_s(M_Z)$:

$$\alpha_s(M_Z) = 0.1161^{+0.0041}_{-0.0048}$$ NLO + 2-loop threshold corrections

$$= 0.1202^{+0.0072}_{-0.0059}$$ NLO

TABLE I: Central values and uncertainties due to different sources for the nine $\alpha_s(p_T)$ results and for the combined $\alpha_s(M_Z)$ result (bottom). All uncertainties are multiplied by a factor of 10^3.

p_T range (GeV)	No. of data points	p_T (GeV)	$\alpha_s(p_T)$	total uncertainty	experimental uncorrelated	experimental correlated	non-perturb. correction	PDF uncertainty	$\mu_{r,f}$ variation
50 - 60	4	54.5	0.1229	$^{+7.6}_{-7.7}$	±0.4	$^{+4.8}_{-4.9}$	$^{+5.8}_{-5.6}$	$^{+0.4}_{-0.6}$	$^{+1.0}_{-1.9}$
60 - 70	4	64.5	0.1204	$^{+6.2}_{-6.3}$	±0.3	$^{+4.1}_{-4.3}$	$^{+4.5}_{-4.3}$	$^{+0.6}_{-0.5}$	$^{+1.3}_{-1.5}$
70 - 80	3	74.5	0.1184	$^{+5.6}_{-5.6}$	±0.3	$^{+3.8}_{-3.9}$	$^{+4.0}_{-3.9}$	$^{+0.6}_{-0.6}$	$^{+1.0}_{-0.9}$
80 - 90	3	84.5	0.1163	$^{+5.1}_{-5.1}$	±0.3	$^{+3.6}_{-3.7}$	$^{+3.5}_{-3.5}$	$^{+0.7}_{-0.7}$	$^{+0.9}_{-0.6}$
90 - 100	2	94.5	0.1142	$^{+5.1}_{-4.9}$	±0.3	$^{+3.5}_{-3.6}$	$^{+3.5}_{-3.3}$	$^{+0.8}_{-0.8}$	$^{+1.1}_{-0.6}$
100 - 110	2	104.5	0.1131	$^{+4.7}_{-4.7}$	±0.2	$^{+3.4}_{-3.5}$	$^{+3.1}_{-3.0}$	$^{+0.8}_{-0.8}$	$^{+1.1}_{-0.6}$
110 - 120	2	114.5	0.1121	$^{+4.2}_{-4.4}$	±0.2	$^{+3.1}_{-3.3}$	$^{+2.5}_{-2.7}$	$^{+0.7}_{-0.8}$	$^{+1.2}_{-0.7}$
120 - 130	1	124.5	0.1102	$^{+4.4}_{-4.4}$	±0.2	$^{+3.2}_{-3.4}$	$^{+2.6}_{-2.6}$	$^{+0.9}_{-0.9}$	$^{+1.4}_{-0.9}$
130 - 145	1	136.5	0.1090	$^{+4.2}_{-4.3}$	±0.3	$^{+3.1}_{-3.4}$	$^{+2.3}_{-2.4}$	$^{+0.9}_{-0.9}$	$^{+1.5}_{-0.9}$
50 - 145	22	M_Z	0.1161	$^{+4.1}_{-4.8}$	±0.1	$^{+3.4}_{-3.3}$	$^{+1.0}_{-1.6}$	$^{+1.1}_{-1.2}$	$^{+2.5}_{-2.9}$

Main correlated uncertainties: JES, pT-resolution, luminosity

α_s: Fit Method

- Minimize χ^2 (used in many PDF fits, D0 dijet angular PRL)

$$\chi^2(\xi,\vec{\epsilon},\vec{\alpha}) = \sum_i \frac{\left[d_i - t_i(\xi,\vec{\alpha})\left(1+\sum_j \delta_{ij}(\epsilon_j)\right)\right]^2}{\sigma^2_{i,\mathrm{stat.}} + \sigma^2_{i,\mathrm{uncorr.}}} + \sum_j \epsilon_j^2 + \sum_k \alpha_k^2$$

→ 23 experimental correlated sources of uncertainty
→ non-perturbative corrections uncertainties
→ PDF uncertainties

Separate treatment for **renormalization and factorization scales** (convention from LEP, HERA):

- perform fits for fixed scale
- repeat for scale factors 2.0, 0.5
- quote differences as 'scale uncertainty'

→ does not assume Gaussian distributed scale uncertainties

Angular distributions: dijet χ (D0)

– Measure χ = exp(|y1 – y2|) in 10 regions of dijet mass with M_{jj} >250 GeV (last bin: >1.1TeV!)
– Good agreement with NLO pQCD(MSTW2008)
– Data are used to set limits on the models of
Quark compositeness: ~ 3 TeV
TeV-1 extra dim. : ~1.6 TeV
ADD extra dim. :~1.3-1.9 TeV (dep. on Ned)

PRL 103, 191803 (2009)

Large excess at small △y is expected in QC and ED models



Small △y Large △y

M>1 TeV:



Jet "Definitions" - Jet Algorithms

Midpoint cone-based algorithm

- Cluster objects based on their proximity in y-ϕ space
- Starting from seeds (calorimeter towers/particles above threshold), find stable cones (kinematic centroid = geometric center).
- Seeds necessary for speed, however source of infrared unsafety.
- In recent QCD studies, we use "Midpoint" algorithm, i.e. look for stable cones from middle points between two adjacent cones
- Stable cones sometime overlap
  → merge cones when p_T overlap > 75%

Infrared unsafety:
soft parton emission changes jet clustering



More advanced algorithm(s) available now, but negligible effects on this measurement.

Jet "Definitions" - Jet Algorithms

k_T algorithm

- Cluster objects in order of increasing their relative transverse momentum (k_T)
  - $d_{ii} = p_{T,i}^2, \quad d_{ij} = \min(p_{T,i}^2, p_{T,j}^2)\ \frac{\Delta R^2}{D^2}$

  until all objects become part of jets
- D parameter controls merging termination and characterizes size of resulting jets



- No issue of splitting/merging. Infrared and collinear safe to all orders of QCD.
- Every object assigned to a jet: concerns about vacuuming up too many particles.
- Successful at LEP & HERA, but relatively new at the hadron colliders
  - More difficult environment (underlying event, multiple *pp* interactions...)

DP Signal variables

Calculate the azimuthal angle for the pair that gives the minimum value of *S*:

$$\Delta S = \Delta\phi\left(\mathrm{p_T}^{\gamma, jet_i},\ \mathrm{p_T}^{jet_j, jet_k}\right)$$

$$S_\phi = \frac{1}{\sqrt{2}}\sqrt{\left(\frac{\Delta\phi(\gamma,i)}{\delta\phi(\gamma,i)}\right)^2 + \left(\frac{\Delta\phi(j,k)}{\delta\phi(j,k)}\right)^2}$$

$$S_{pT} = \frac{1}{\sqrt{2}}\sqrt{\left(\frac{|\vec{P}_T(\gamma,i)|}{\delta P_T(\gamma,i)}\right)^2 + \left(\frac{|\vec{P}_T(j,k)|}{\delta P_T(j,k)}\right)^2}$$





MINIMUM BIAS – HYPERON PRODUCTION (CDF)

→ Strange particle production can reveal mechanisms from the collision.







→ Cross sections are measured in pT bins, accessing previously unexplored high pT regions.

→ Cross sections are also measured in different multiplicity regions.

http://www-cdf.fnal.gov/physics/new/qcd/hyperons_10/hyperon_public_note.pdf

MPI, experimental tests

Jet direction

Jet pedestal effect





Effective parton Luminosity:

$$L_{eff}(\Delta)=\int D(r)D(r')dV_{overlap}$$

CDF (Run 2)



- Presence of high pT 1st interaction biases events towards smaller p-pbar impact parameters and hence leads to a higher additional activity but saturates at σ(pT_jet) ≪ σ_nd ("nd" = non-diffractive).
- The height of the pedestal depends on the overlap, i.e. on the parton matter distribution function.

b-bbar Dijet Production (CDF)

Preliminary

- Preliminary cross section results with L = 260 pb-1
- jet pT>35 and 32 GeV, |eta|<1.2
- The purity of b-bbar events is calculated using SVT track mass; purities in the mass/$\triangle\Phi$ bins are 75-90%
- Comparison with Pythia (tune A), Herwig+Jimmy and MC@NLO+Jimmy:

Data: σ = 5664 ± 168(stat) ± 1270 (syst) pb
Pythia: σ = 5136 ± 52(stat)
Herwig: σ = 5296 ± 98(stat)
MC@NLO: σ = 5421 ± 105(stat)

- Tested: lead.jet pT, dijet mass, $\triangle\Phi$; good agreement
- Discrepancy with MC gen. predictions at small $\triangle\Phi$.







Z+jets production. Δϕ(Z,jet)

First measurement of $\Delta\phi$(Z, jet) !

- $Z \to \mu\mu$, $|y_\mu|<1.7$, $p_{TZ} > 25$ GeV
- jet $p_T>20$ GeV, |jet y| < 2.8

PLB 682, 370 (2010)



PYTHIA p_T ordered
- new "Perugia" tune
- MRST07 LO* PDF
PYTHIA Q^2 ordered
HERWIG

ALPGEN + PYTHIA p_T
ALPGEN + PYTHIA Q^2
ALPGEN + HERWIG

- Sherpa describes Δϕ(Z,jet) shape very well (but a normalization issue)
- Small values of Δϕ are excluded from MCFM due to significant non-perturbative contributions

Z+jets production. Δy(Z,jet)

First measurement of Δy(Z, jet) !

- $Z \to \mu\mu$, $|y_\mu|<1.7$, $p_{TZ} > 25$ GeV
- jet $p_T>20$ GeV, |jet y| < 2.8



PYTHIA p_T ordered
- new "Perugia" tune
- MRST07 LO* PDF
PYTHIA Q^2 ordered
HERWIG

ALPGEN + PYTHIA p_T
ALPGEN + PYTHIA Q^2
ALPGEN + HERWIG

Sherpa, NLO describe Δy

Three jet mass: χ^2 test (D0)