

# Exploring QCD with COMPASS-II

C. Quintans, LIP-Lisbon

12<sup>th</sup> April 2011, DIS 2011




Co-financed by:




FCT Fundação para a Ciência e a Tecnologia  
MINISTÉRIO DA CIÊNCIA, TECNOLOGIA E ENSINO SUPERIOR Portugal

Fundação Luso-Americana



- ◆ Unpolarized Deep Inelastic Scattering and the strange quark PDF
- ◆ Deeply Virtual Compton Scattering and Generalized Parton Distributions
- ◆ Drell-Yan and Transverse Momentum Dependent PDFs
- ◆ Primakoff reaction as a test of Chiral Perturbation Theory
- ◆ Summary




## ***A bit of history***

---

COMPASS has been taking data at the SPS/CERN since 2002, with main goals:

- ◆ Helicity PDF  $g_1$  from double polarized DIS, and flavor separation
- ◆ Gluon polarization  $\Delta G/G$  from double polarized DIS
- ◆ Transversity PDF  $h_1$  in transversely polarized SIDIS
- ➔ Naturally polarized  $\mu^+$  beam (@160 GeV/c), on  ${}^6\text{LiD}$  and  $\text{NH}_3$  transversely or longitudinally polarized targets.
  
- ◆ Hadron spectroscopy: searches of exotics, hybrids and glueballs
- ◆ Pion polarizabilities
- ➔ Unpolarized hadron beams (@190 GeV/c) on several unpolarized targets (Liquid  $\text{H}_2$ , Pb, Ni, Cu and W)



# Studying QCD in COMPASS...


---

Thanks to a multi-purpose spectrometer, and a versatile beam line, COMPASS has been doing extensive QCD studies.

The **nucleon spin puzzle**, that has been intriguing us since EMC times, comes now with new clues:


- ◆ The quarks spin is responsible for only  $\approx 30\%$  of the nucleon spin
- ◆ The gluons spin does not contribute much, at least in the x-region scanned by COMPASS ( $x_g \approx 0.1$ )
- ◆ What about orbital angular momentum contributions from quarks and gluons?

In LO 3 PDFs are needed for a full description of the nucleon: unpolarized, helicity and transversity – all investigated in COMPASS.




But the 1-dimensional picture of the nucleon is now to be improved:


- ◆ quarks flavor matters: improve namely the strange PDF;
- ◆ 3-dimensional description, via GPDs;
- ◆ account for intrinsic transverse momentum of partons, via TMD PDFs;
- ◆ ...and low energies QCD: a test of Chiral Perturbation Theory.




# COMPASS Experiment




*Polarized target:  ${}^6\text{LiD}$  or  $\text{NH}_3$*




$\mu^\pm, \pi^-$  or  $p$  beam






# Deep Inelastic Scattering




DIS is used since long to access the **parton distributions**  $q(x)$ .

Flavor separation: identify one outgoing hadron, besides  $\mu'$  – SIDIS.


While the unpolarized light quark PDFs are well constrained, the **strange quark distribution** has still a large uncertainty:






# Unpolarized SIDIS

Unpolarized SIDIS measurement, with PID and charge separation from the COMPASS RICH and Calorimeters:  $K^\pm$ ,  $K^0$ ,  $\pi^\pm$ ,  $\pi^0$ ,  $\Lambda$ ,...



- ◆ Improve on **unpolarized PDFs**, namely  $s(x)$
- ◆ Improve on **fragmentation functions**, namely kaon FF
- ◆ LO analysis of COMPASS data alone
- ◆ Scan PDFs and FFs on  $(x, z, \dots)$ , towards a NLO global analysis


Measurement in parallel with the GPDs program, using a long liquid  $H_2$  target.




# Generalized Parton Distributions

**GPDs:** a 3D picture of the nucleon, by adding information about the transverse distance of the constituent quark.

They allow to access information on the quarks orbital angular momentum.




- ◆ 4 GPDs:  $H$ ,  $E$ ,  $\tilde{H}$  and  $\tilde{E}$ , for each quark flavor and gluons.
- ◆ They depend on 3 variables:  $x$ ,  $\xi = \frac{x_B}{2-x_B}$  and  $t$ .



# DVCS in COMPASS

Transverse imaging (tomography) of the nucleon from **Deeply Virtual Compton Scattering** (DVCS):  $\mu p \rightarrow \mu' p \gamma$




$$\frac{d\sigma}{dt} \approx e^{-Bt} \quad B \approx \langle r_{\perp}^2 \rangle / 2$$


- ◆ Main priority is DVCS.
- ◆ Deeply Virtual Meson Production (DVMP):  $\mu p \rightarrow \mu' p \rho$  will be studied in parallel.
- ◆ 2 competing processes: DVCS and Bethe-Heitler
  - ★ Low  $x_B$ : BH;
  - ★ High  $x_B$ : DVCS;
  - ★ intermediate  $x_B$ : interference DVCS-BH.
- ◆ BH is well-known: used as reference process.



**Phase 1:**  $\mu^{+\downarrow}$  and  $\mu^{-\uparrow}$  beams off a 2.5m unpolarized liquid  $H_2$  target  $\Rightarrow$  **GPD H.**  
**Phase 2:**  $\mu^{+\downarrow}$  and  $\mu^{-\uparrow}$  beams off a transversely polarized  $NH_3$  target  $\Rightarrow$  **GPD E.**



# DVCS: phase-space and goals




Measure the modulations given by the **sum** and **difference** of "spin and charge" dependent DVCS cross-sections, whose amplitudes are proportional to  $\mathcal{I}m(F_1 H)$  and  $\mathcal{R}e(F_1 H)$

$$\mathcal{S}_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) + d\sigma(\mu^{-\uparrow})$$

$$\mathcal{D}_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) - d\sigma(\mu^{-\uparrow})$$

COMPASS will cover an unexplored region of the phase-space, the intermediate  $x_B$ .



The measurement requires a **recoil proton detector**, and large coverage **electromagnetic calorimetry**.




# DVCS: Projections


Phase 1:  $B = \langle r_{\perp}^2 \rangle / 2$

$$D_{CS,U} \equiv d\sigma(\mu^{+\downarrow}) - d\sigma(\mu^{-\uparrow})$$




Phase 2: Azimuthal asymmetries





**TMDs**: dynamic picture of the nucleon, by considering also the partons intrinsic transverse momentum  $k_T$ .

They also allow to access information on the quarks orbital angular momentum.



In COMPASS TMD PDFs can be studied in **2 complementary ways**:


- ◆ **Semi-inclusive DIS**







  - ↪ polarized muon beam on unpolarized/transversely polarized target;

- ◆ **Drell-Yan process**

  - ↪ pion beam on unpolarized/transversely polarized target.

# TMDs approach



		NUCLEON		
		unpolarized	longitudinally pol.	transversely pol.
QUARK	unpolarized	$f_1$  number density		$f_{1T}^\perp$  Sivers
	longitudinally pol.		$g_{1L}$  helicity	$g_{1T}$ 
	transversely pol.	$h_1^\perp$  Boer-Mulders	$h_{1L}^\perp$ 	$h_1$  transversity $h_{1T}^\perp$  pretzelosity


In LO and considering the quarks  $k_T$ , 8 PDFs are needed to fully describe the nucleon.

The TMDs approach is valid when

$$Q \gg q_T \gtrsim \Lambda_{QCD}.$$

TMDs are accessed by measuring azimuthal asymmetries.


## SIDIS



The spin asymmetry is proportional to PDF  $\otimes$  FF:


$$A_{Sivers} \propto \frac{\sum_q e_q^2 f_{1T}^{\perp(1)}(x) D_q^h(z)}{\sum_q e_q^2 f_1(x) D_q^h(z)}$$

## DY




The spin asymmetry is proportional to PDF<sup>b</sup>  $\otimes$  PDF<sup>t</sup>. If unpolarized beam and transversely polarized target:

$$A_{Sivers} \propto 2 \frac{\sum_q e_q^2 \bar{f}_{1q}(x_1) f_{1Tq}^{\perp(1)}(x_2)}{\sum_q e_q^2 f_{1q}(x_1) f_{1q}(x_2)}$$




# Boer-Mulders and Sivers in SIDIS

We look at the azimuthal modulations of produced hadrons,  $\phi$  being the angle of the hadron in the  $\gamma^*$ -nucleon reference frame.



2004-2006 data on deuteron target: non-zero **Boer-Mulders asymmetry** ( $A_{LU}^{\cos 2\phi}$ ).

The Boer-Mulders TMD in the proton can be measured in parallel with DVCS.



The **Sivers asymmetry** ( $A_{LT}^{\sin \phi_S}$ ) was measured in COMPASS in deuteron and proton targets.


Positive asymmetry for  $h^+$  on proton, but effect seen in HERMES is a factor 2 larger.

As Sivers and Boer-Mulders are T-odd PDFs:

$$f_{1T}^\perp(DY) = -f_{1T}^\perp(SIDIS)$$


$$h_1^\perp(DY) = -h_1^\perp(SIDIS)$$

↪ A crucial test of non-perturbative QCD and of TMDs approach.




# TMDs in Drell-Yan

Drell-Yan events from  $\pi^-$  @190 GeV/c collisions on a  $NH_3$  target transversely polarized  $\Rightarrow$  access 4 azimuthal modulations related to **Boer-Mulders**, **Sivers**, **pretzelosity** and **transversity** PDFs.







- ◆ All 4 asymmetries predicted to be sizable in the valence quarks region.
- ◆ COMPASS acceptance is very favorable:  $x_p > 0.05$ .
- ◆ Study Drell-Yan in  $4 < M_{\mu\mu} < 9$  GeV/c<sup>2</sup>, where background is negligible.
- ◆ TMD approach validity guaranteed by  $M_{\mu\mu} \gg p_T^{\mu\mu} \approx 1$  GeV.

The measurement requires a **hadron absorber** downstream the target, and a **dimuon trigger** system.




# Drell-Yan: projections

DY 4. – 9. GeV/c<sup>2</sup>




Assuming 2 years of data-taking, it will be possible to check the sign change between SIDIS and DY, and test PDF predictions.




**Chiral Perturbation Theory** predicts the strong interaction dynamics of Goldstone bosons.

↪ the internal structure of the pion is revealed by its response in presence of an electromagnetic field, i.e. **pion polarizabilities**.




Primakoff



Inverse Compton

Studying the  $\pi$  induced Primakoff reaction and the embedded inverse Compton scattering, one can measure the  $\pi$  polarizabilities and check ChPT predictions.




# Pion polarizabilities

$$\frac{d\sigma_{\pi\gamma}}{d\Omega_{cm}} = \left[ \frac{d\sigma_{\pi\gamma}}{d\Omega_{cm}} \right]_{point-like} + C \frac{s - m_\pi^2}{s^2} \mathcal{P}(\alpha_\pi, \beta_\pi)$$

Polarizability:

$$\begin{aligned} \mathcal{P} = & (1 - \cos \theta_{cm})^2 (\alpha_\pi - \beta_\pi) \\ & + (1 + \cos \theta_{cm})^2 (\alpha_\pi + \beta_\pi) \frac{s^2}{m_\pi^4} \\ & + (1 - \cos \theta_{cm})^3 (\alpha_2 - \beta_2) \frac{(s - m_\pi^2)^2}{24s} \end{aligned}$$




In 120 days (90 with $\pi$ , 30 with $\mu$ beams)	$\alpha_\pi - \beta_\pi$ ( $10^{-4} \text{ fm}^3$ )	$\alpha_\pi + \beta_\pi$ ( $10^{-4} \text{ fm}^3$ )	$\alpha_2 - \beta_2$ ( $10^{-4} \text{ fm}^5$ )
2-loop ChPT prediction	$5.7 \pm 1.0$	$0.16 \pm 0.10$	16
<b>COMPASS sensitivity</b>	<b><math>\pm 0.66</math></b>	<b><math>\pm 0.025</math></b>	<b><math>\pm 1.94</math></b>

Measurement with muon beam will allow to cross-check systematics.

Up to now, experiments measured  $\alpha_\pi - \beta_\pi$  from  $4$  to  $14 \times 10^{-4} \text{ fm}^3$ .

In addition COMPASS will measure for the first time **kaon polarizabilities**.



# ***Timeline and Conclusions***

---

COMPASS **new proposal** (May 2010), with physics program including:

- ◆ Unpolarized SIDIS measurements to access the strange quark PDF, FFs and TMD PDFs with unprecedented precision;
- ◆ DVCS and DVMP for GPDs studies in a phase-space region not yet covered by other past or near-future experiments;
- ◆ First polarized Drell-Yan experiment for TMDs studies;
- ◆ Primakoff reactions for pion polarizabilities (improved accuracy) and first measurement of kaon polarizabilities.

↪ at least 5 years of data-taking, which can start from 2012.

Program approved by the SPSC/CERN for a first period of 3 years.

COMPASS will continue playing a major role in QCD physics for the next 5 to 10 years.

**Stay tuned!**